



Veterinærinstituttet
Norwegian Veterinary Institute

⋮ Annual Report

The surveillance programme for *enzootic bovine leukosis* (EBL) in Norway 2019



Comissioned by



The surveillance programme for *enzootic bovine leukosis* (EBL) in Norway 2019

Content

Summary	3
Introduction	3
Aim	3
Materials and methods	3
Results	4
Discussion	5
References	5

Authors

Thea Blystad Klem, Johan Åkerstedt and Siv Klevar

Commissioned by

Norwegian Food Safety Authority



ISSN 1894-5678

© Norwegian Veterinary Institute 2020

Design Cover: Reine Linjer

Photo front page: Colourbox

Summary

All milk and blood samples tested in 2019 were negative for antibodies against bovine leukaemia virus (BLV).

Introduction

Enzootic bovine leukosis (EBL) is caused by bovine leukaemia virus (BLV), in the genus retrovirus. Most infections are subclinical. Approximately one third of infected cattle older than three years of age develop persistent lymphocytosis. A smaller proportion of animals develop lymphosarcomas in various internal organs. EBL is classified as a list B disease in Norway and is notifiable to the Office International des Epizooties. The disease had never been reported in Norway until antibodies against BLV were detected in eight dairy herds in samples collected through the surveillance programme in 1995 (1). No new herds have tested positive since 1997 (2), except a single positive bulk milk sample in one herd in 2002, and one blood sample from one of the cows in that herd. After extensive follow up, it was concluded that the positive antibody results were due to false positive reactions (3). Free status from EBL was granted to Norway by the EFTA Surveillance Authority in 2007.

The Norwegian Food Safety Authority is responsible for carrying out the surveillance programme for EBL. The Norwegian Veterinary Institute is in charge of planning the programme, collecting the bulk milk samples from the dairies, and performing the tests. Official inspectors from the Norwegian Food Safety Authority collected the blood samples from beef cattle at slaughterhouses.

Aim

The aim of the surveillance programme for EBL is to document freedom from the infection in Norway according to Council Directive 64/432/EEC as amended and to contribute to the maintenance of this favourable situation.

Materials and methods

The surveillance programme included both dairy and beef herds. The target population of dairy herds consisted of all cattle herds delivering milk to dairies during the sampling period. The target population of beef herds was all herds delivering cattle to slaughter in 2019. Sixteen and a half per cent of the dairy herds were randomly selected for sampling. Bulk milk samples were provided by the dairies. From the beef herds, individual blood samples from animals older than 24 months were collected at 18 slaughterhouses, with a maximum of five animals per herd and day of sampling. Bulk milk samples from 1071 dairy herds were collected and tested. A total of 4124 individual blood samples from 1328 beef herds were received for analyses in pools (n = 1685). The sampled herds represented approximately 17.9% of the Norwegian cattle herds (Table 1).

Table 1. Numbers of dairy herds and beef herds and herds sampled within the frame of the Norwegian surveillance programme for EBL in 2019.

Herd category	Cattle herds (total no. ¹)	Sampled herds (no. ²)	Sampled herds (%)
Dairy herds ³	7 600	1 071	14.1
Beef herds ⁴	5 503	1 328	24.1
Total	13 352	2 389	17.9

¹Based on data from the Register of production subsidies as of 1 March 2019.

²Combined beef cattle and dairy farms could be sampled under both herd categories. Number of unique farms is given as total number of sampled herds.

³Cattle herds delivering milk to dairies.

⁴Sampling performed at slaughterhouses

Blood samples (pooled or individual samples) were examined using an indirect ELISA SVANOVIR® BLV gp51-Ab from Boehringer Ingelheim Svanova (Uppsala, Sweden) and/or using a competitive ELISA kit for Enzootic Bovine Leukosis Virus ID Screen® BLV Competition (ID.Vet, Montpellier, France). In case of positive or inconclusive results in pooled blood samples, individual samples were re-tested. Individual samples with inconclusive or positive results were re-tested in duplicates using the Enzootic Bovine Leukosis Virus ID Screen® BLV Competition (ID.Vet, Montpellier, France).

Bulk milk samples were analysed using an indirect ELISA, IDEXX Leukosis Milk Screening Ab test (IDEXX Laboratories, Maine, USA). Inconclusive and positive reactors were retested in duplicates using the same method. In case of doubtful reactors, new blood samples/bulk milk samples from the suspected animals or herd, were requested and tested. All samples were analysed at the Norwegian Veterinary Institute.

Results

All bulk milk samples and blood samples tested in 2019 were negative for antibodies against BLV. Table 2 shows the results of the testing during the period from 1995 to 2019.

Table 2. Numbers of samples and positive results of antibody testing in the surveillance programme for EBL in the Norwegian cattle population during the period 1995-2019.

Year	Dairy herds	Beef herds		No. of positive samples
	No. of herds sampled	No. of herds sampled ¹	No. of individuals tested ²	
1995	25 131	1 532	9 354	8 (bulk milk)
1996	2 278	303	1 523	1 (bulk milk)
1997	26 903	2 214	16 741	0
1998	23 581	2 191	17 095	0
1999	19 933	2 382	18 274	0
2000	1 590	340	2 892	0
2001	2 564	434	3 453	0
2002	2 308	462	3 693	1 (bulk milk)
2003	1 845	449	3 901	0
2004	1 573	402	3 364	0
2005	1 919	484	4 766	0
2006	1 673	479	4 624	0
2007	1 575	412	4 241	0
2008	1 422	444	4 616	0
2009	1 315	435	5 038	0
2010	1 265	507	4 020	0
2011	1 226	1 278	4 758	0
2012	1 189	1 178	4 306	0
2013	1 042	1 167	4 079	0
2014	1 489	935	4 132	0
2015	1 176	1 206	3 704	0
2016	1 180	1 337	4 241	0
2017	1 107	1 448	4 285	0
2018	1 131	1 341	4 153	0
2019	1 071	1 328	4 124	0

¹Sampling performed in the herds prior to 2011.

²A small number of blood samples collected at slaughterhouses could originate from dairy herds.

Discussion

The requirement from the EU for granting an EBL-free status is that the herd prevalence must be lower than 0.2%, which represents a maximum of 26 herds out of the total number of 13 352 herds.

No new cases have been reported after 1997, and the continuous surveillance since 1995 shows that the Norwegian cattle population is free from EBL according to the requirements (2, 3). Initially, all cattle herds were tested annually. Since 2000, a minimum of 10% of dairy and beef cattle herds have been tested each year. Using scenario tree modelling, the probability of freedom from EBL in Norway at the end of 2014 was calculated to 99.0% (5). The results of the surveillance programme from 2019 support that the Norwegian cattle population is free of EBL.

Together with the possible isolation period of six months and the testing protocol for imported animals, the surveillance programme for EBL should be an effective means to detect introduction of new infection.

References

1. Tharaldsen J, Ødegaard Ø, Krogsrud J. Smittsom storfeleukose diagnostisert i Norge [Contagious bovine leukosis diagnosed in Norway, No]. *Nor Vet Tidsskr* 1996; 108: 550.
2. Åkerstedt J, Skar K, Hopp P, Klem TB. The surveillance programme for enzootic bovine leucosis (EBL) in Norway 2018. Surveillance Programmes for terrestrial and aquatic animals in Norway. Annual report 2018. Oslo: Norwegian Veterinary Institute 2019.
3. Nyberg O, Tharaldsen J, Heier BT. The surveillance and control programme for enzootic bovine leukosis (EBL) in Norway. In: Mørk T, Hellberg H (editors). Surveillance and control programmes for terrestrial and aquatic animals in Norway. Annual report 2003. Oslo: National Veterinary Institute; 2004. p. 57-62.
4. Klintevall K, Näslund K, Svedlund G, Hajdu L, Linde N, Klingeborn B. Evaluation of an indirect ELISA for the detection of antibodies to bovine leukemia virus in milk and serum. *J Virol Methods* 1991; 33: 319-33.
5. Jonsson ME, Kristoffersen AB, Åkerstedt J, Hopp P, Mørk T, Sjurseth SK, Norström M. Evaluation of the surveillance sensitivity of the official Enzootic bovine leukosis programme in Norway. Annual Meeting of the Society for Veterinary Epidemiology and Preventive Medicine, Elsinore, Denmark, 16-18 March 2016. Poster.

Frisk fisk



Sunne dyr



Trygg mat



Faglig ambisiøs, fremtidsrettet og samspillende - for én helse!



Veterinærinstituttet
Norwegian Veterinary Institute

Oslo

Trondheim

Sandnes

Bergen

Harstad

Tromsø

postmottak@vetinst.no
www.vetinst.no