


The surveillance and control programme for viral haemorrhagic septicaemia (VHS) and infectious haematopoietic necrosis (IHN) in Norway

Hege Hellberg

Irene Ørpetveit

Britt Bang Jensen



*Scientific editors Hege Hellberg and Ståle Sviland
Norwegian Veterinary Institute*



**Surveillance and control programmes for terrestrial and aquatic animals in Norway.
Annual report 2010**

Editors

Ståle Sviland and Hege Hellberg

Publisher

Norwegian Veterinary Institute
PO Box 750 Sentrum
N-0106 Oslo
Norway

Fax: + 47 23 21 60 01

Tel: + 47 23 21 60 00

E-mail: postmottak@vetinst.no

www.vetinst.no

Front page photo: Processed from Colourbox

ISSN 1503-1454

Example of citation:

Hellberg H, Ørpetveit I, Bang Jensen B. The surveillance and control programme for viral haemorrhagic septicaemia (VHS) and infectious haematopoietic necrosis (IHN) in Norway. In: Sviland S, Hellberg H (editors). Surveillance and control programmes for terrestrial and aquatic animals in Norway. Annual report 2010. Oslo: Norwegian Veterinary Institute; 2011. ISSN 1503-1454.

© Norwegian Veterinary Institute

Any use of the present data should include specific reference to this report.

The surveillance and control programme for viral haemorrhagic septicaemia (VHS) and infectious haematopoietic necrosis (IHN) in Norway

Hege Hellberg, Irene Ørpetveit, Britt Bang Jensen

*The surveillance programme was amended in 2009 and sampling was limited to rainbow trout (*Oncorhynchus mykiss*). VHS virus (VHSV) and IHN virus (IHNV) was not detected at any of the sites tested in the 2010 surveillance. Only farms in the approved zone are included in the surveillance programme.*

Introduction

VHS and IHN are two important diseases in salmonid fish caused by rhabdovirus infections (1). VHS occurs in several fish species in both freshwater and seawater, but has most frequently been recorded in farmed rainbow trout. The VHS outbreak in the Great Lakes in North America continues to cause large losses among several wild fish species (2, 3). VHS was diagnosed in farmed rainbow trout in Norway in 2007 (4). Outbreaks of IHN have resulted in significant economic losses in farmed rainbow trout and salmon in North America and Europe, and the disease has also had an impact on wild populations of Pacific salmon. IHN has never been diagnosed in Norway. For more information on fish rhabdoviruses:

<http://www.vetinst.no/nor/Faktabank/Alle-faktaark/Viral-hemoragisk-septikemi>

<http://www.vetinst.no/nor/Faktabank/Alle-faktaark/Infeksioes-Hematopoetisk-Nekrose>

<http://www.vetinst.no/eng/Research/Publications/Fish-Health-Report>

<http://www.vetinst.no/eng/Research/Publications/Surveillance-and-Control-Programs-annual-reports>

Aim

The aim of the programme is to document the absence of VHSV and IHNV in fish farms within the approved zone in order to maintain Norway's VHS and IHN free status.

Materials and methods

Sampling

Sampling and inspection is carried out by the Norwegian Food Safety Authority District Offices according to Directive 2006/88/EC and Decision 2001/183/EC (5, 6). The virological analyses are performed by the Norwegian Veterinary Institute. Thirty fish are sampled from each site and, depending on fish size, samples from five or ten individuals are pooled on transport medium.

Analysis

The samples are processed according to Decision 2001/183/EC (6) and recommendations from EU reference laboratory for fish diseases in Århus, Denmark. Following homogenisation and low speed centrifugation, the resulting supernatant is incubated with a polyclonal antibody against infectious pancreatic necrosis virus (IPNV) and analysed in cell culture.

Results

In 2010, a total of 115 pooled samples (1,065 individual fish) from 36 sites were examined (Tables 1 and 2, Figure 1). All samples were negative for IHN and VHS.

Table 1. Different categories of fish analysed for VHSV and IHN in 2010

	Fry - smolt		On-growing		Brood fish		Total	
	No. sites	No. of pooled samples	No. sites	No. of pooled samples	No. sites	No. of pooled samples	No. sites	No. of pooled samples
Rainbow trout (<i>Oncorhynchus mykiss</i>)	10	38	25	74	-	-	35	112
Brown trout (<i>Salmo trutta</i> L.)	-	-	1	3	-	-	1	3
Total	10	38	26	77	-	-	36	115

Table 2. Number of farms and species analysed for VHSV and IHN during the time period 1995-2010

Farm types	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
<i>Per production type</i>																
Smolt producing farms	71	169	162	30	27	45	30	32	54	51	125	104	140	84	17	10
On-growing farms	207	340	346	478	527	447	508	414	429	303	280	276	282	315	32	26
Brood stock farms				2	3	7	7	14	2	9	14	15	12	48	1	
<i>Per species</i>																
Farms with Atlantic salmon	225	425	392	417	462	382	408	372	387	295	345	316	359	376	0	
Farms with rainbow trout	31	63	69	66	62	83	93	61	74	48	61	49	54	48	49	35
Farms with brown trout	15	13	38	21	27	28	24	23	24	21	8	24	13	17	1	1
Farms with char	1	7	6	5	4	10	8	9	9	5	7	8	5	8		
Farms with turbot	6	1	1		1	1	4		1	1		1		1		
Farms with sea trout				2	3	2	4	1	2	2	2	2	2			
Farms with brook trout				2		1	1	2	1	2						
Farms with relict Atlantic salmon				1						1						
Farms with cod													5			
Total	278	509	506	510	554	494	534	468	498	375	417	392	434	444	50	36

* During the period 1995-2008, cultivation facilities for wild stocks were categorized as brood stock farms.

Discussion

In 1994, Norway obtained disease free status for VHS and IHN (7). The 2007 VHS outbreak in rainbow trout caused a temporary suspension of the disease free status. Measures to eliminate the disease and prevent its spread were immediately taken by the Food Safety Authority. In May 2008, Norway regained its VHS free status, with the exception of the VHS outbreak area (8). The surveillance programme only covers the approved zone.

In 2009, 3 pooled samples from 1 site were rejected, compared to 1 pooled sample from 1 site in 2008. In 2010, 3 pooled samples from 1 site were rejected, the same number as in 2009.

The isolation of SAV (the virus causing pancreas disease - PD) in samples received for surveillance of VHSV and IHNV continues to represent a problem for detection of the two rhabdoviruses. Presently, neutralising antibodies against SAV are not available. Also, it is not known whether replication of SAV will inhibit replication of VHSV in BF-2 cells. Therefore, tissue homogenates of SAV-positive samples are always examined for VHSV by real-time RT-PCR to ensure the absence of this virus. The SAV-positive sites are located in the region where PD is considered endemic.

The farms in the VHS outbreak area have been subjected to targeted surveillance with a view to restoration of approved zone status. In 2010 a total of 165 pooled samples (1515 individual fish) from 7 sites have been analysed. All samples were negative for IHNV and VHSV.

Conclusion

Based on the examinations carried out in the surveillance and control programme for VHS and IHN at the Norwegian Veterinary Institute in 2010, no confirmed or suspected cases of VHSV or IHNV were registered within the approved zone. Prior to the 2007 VHS outbreak, none of the involved sites had tested positive in the surveillance programme. The surveillance programme was evaluated in 2010 and the programme will be amended in 2011. For more information on the evaluation of the programme:

<http://www.vetinst.no/nor/Forskning/Publikasjoner/Rapportserie>

References

1. Anonymous. Diseases of Fish. In: Manual of diagnostic tests for aquatic animals 2010. Part 2, Paris: Office International des Epizooties; 2010.
2. Elsayed E, Faisal M, Thomas M, Whelan G, Batts W, Winton J. Isolation of viral haemorrhagic septicaemia virus from muskellunge, *Esox masquinongy* (Mitchill), in Lake St. Clair, Michigan, USA reveals a new sublineage of the North American genotype. J Fish Dis 2006; 29: 611-19.
3. Lumsden JS, Morrison B, Yason C, Russell S, Young K, Yazdanpanah A, Huber P, Al-Hussiney L, Stone D, Way K. Mortality event in freshwater drum *Aplodinotus grunniens* from Lake Ontario, Canada, associated with viral haemorrhagic septicaemia virus, Type IV. Dis Aquat Org. 2007; 76: 99-111.
4. Dale OB, Ørpetveit I, Lyngstad TM, Kahns S, Skall HF, Olesen NJ, Dannevig BH. Outbreak of viral haemorrhagic septicaemia (VHS) in seawater-farmed rainbow trout in Norway caused by VHS virus genotype III. Dis Aquat Org. 2009; 85: 93-103.
5. Council Directive 2006/88/EC of 24 October 2006 on animal health requirements for aquaculture animals and products thereof, and on the prevention and control of certain diseases in aquatic animals.
6. Commission Decision 2001/183/EC of 22 February 2001 laying down the sampling plans and diagnostic methods for the detection and confirmation of certain fish diseases.
7. EFTA Surveillance Authority Decision No. 71/94/COL of June 1994.
8. EFTA Surveillance Authority Decision No. 302/08/COL of May 2008.

The Norwegian Veterinary Institute (NVI) is a nationwide research institute in the fields of animal health, fish health, and food safety. The primary mission of the NVI is to give research-based independent advisory support to ministries and governing authorities. Preparedness, diagnostics, surveillance, reference functions, risk assessments, and advisory and educational functions are the most important areas of operation.

The Norwegian Veterinary Institute has its main laboratory in Oslo, with regional laboratories in Sandnes, Bergen, Trondheim, Harstad og Tromsø, with about 360 employees in total.

www.vetinst.no



Veterinærinstituttet
Norwegian Veterinary Institute

The Norwegian Food Safety Authority (NFSA) is a governmental body whose aim is to ensure through regulations and controls that food and drinking water are as safe and healthy as possible for consumers and to promote plant, fish and animal health and ethical farming of fish and animals. We encourage environmentally friendly production and we also regulate and control cosmetics, veterinary medicines and animal health personnel. The NFSA drafts and provides information on legislation, performs risk-based inspections, monitors food safety, plant, fish and animal health, draws up contingency plans and provides updates on developments in our field of competence.

The NFSA comprises three administrative levels, and has some 1300 employees.

The NFSA advises and reports to the Ministry of Agriculture and Food, the Ministry of Fisheries and Coastal Affairs and the Ministry of Health and Care Services.

www.mattilsynet.no

