



## The surveillance programme for scrapie in Norway 2023



REPORT 27/2024

## The surveillance programme for scrapie in Norway 2023

### Authors

Jorunn Mork, Sylvie Lafond Benestad, Lars Austbø, Torfinn Moldal, Petter Hopp

### Suggested citation

Mork, Jorunn, Benestad, Sylvie Lafond, Austbø, Lars, Moldal, Torfinn, Hopp, Petter. The surveillance programme for scrapie in Norway 2023. Surveillance program report. Veterinærinstituttet 2024. © Norwegian Veterinary Institute, copy permitted with citation

### Quality controlled by

Merete Hofshagen, Director of Animal Health, Animal Welfare and Food Safety, Norwegian Veterinary Institute

### Published

2024 on [www.vetinst.no](http://www.vetinst.no)

ISSN 1890-3290 (electronic edition)

© Norwegian Veterinary Institute 2024

### Commissioned by / In collaboration with

Norwegian Food Safety Authority



### Colophon

Cover design: Reine Linjer

Cover photo: Colourbox

[www.vetinst.no](http://www.vetinst.no)

# Content

<b>Summary</b> .....	<b>3</b>
<b>Introduction</b> .....	<b>3</b>
<b>Aims</b> .....	<b>4</b>
<b>Materials and methods</b> .....	<b>4</b>
Animals with clinical signs consistent with scrapie .....	5
Surveillance of fallen stock.....	5
Abattoir surveillance .....	5
Laboratory examination procedures .....	5
PrP genotyping .....	5
Prevalence .....	6
<b>Results</b> .....	<b>6</b>
Sheep .....	6
Goat .....	7
<b>Discussion</b> .....	<b>8</b>
<b>Acknowledgment</b> .....	<b>9</b>
<b>References</b> .....	<b>9</b>


## Summary

In 2023, 19,188 sheep and 568 goats were examined for prion protein scrapie. Seven sheep from seven flocks were positive for scrapie Nor98. All the goats were negative for prion protein scrapie. In the period between 1981 and 2023, scrapie was diagnosed in 278 sheep flocks, of which 221 was Nor98. One goat herd was diagnosed with Nor98.

## Introduction

Scrapie was first diagnosed in indigenous Norwegian sheep in 1981. Increasing numbers of scrapie-infected flocks were identified in the 1990s, culminating with 31 detected flocks in 1996 (Figure 1).

Based on these findings the Norwegian scrapie surveillance programme was launched in 1997 (1).



**Figure 1:** Annual number of sheep flocks and goat herds diagnosed with classical scrapie and Nor98 scrapie during the time period 1980-2023. Before 1998 the cases were not classified according to type of scrapie, but the majority of the scrapie cases, if not all, are considered to have been the classical type.

In 1998, a new type of scrapie, Nor98 scrapie, was identified in Norway. The diagnosis of Nor98 scrapie is verified by Western blot. Nor98 scrapie differs from classical scrapie in several aspects, including the Western blot profile, the distribution of protease resistant prion protein (PrP<sup>Sc</sup>) in the brain, and absence of detectable PrP<sup>Sc</sup> in lymphoid tissues (2). The main clinical sign observed in Nor98 scrapie cases has been ataxia. The PrP genotype distribution among Nor98 scrapie cases differs markedly from that of the cases with classical scrapie (3). Nor98 scrapie is considered to be not contagious or only contagious to a low degree between live animals under field conditions, and is hypothesised to have a spontaneous aetiology.

Scrapie has been a notifiable disease in Norway since 1965. The control measures have involved destruction of all sheep in affected flocks and in close contact flocks. Since 2004, the destruction of all sheep have been performed only for flocks with classical scrapie.

The Norwegian Food Safety Authority (NFSA) is responsible for carrying out the surveillance programme for scrapie. The samples from small ruminants with clinical signs consistent with scrapie and sheep at the abattoirs are collected by inspectors from the NFSA. The NFSA carries out inspections of sheep flocks and goat herds. Since 2019, the company Biosirk, a company collecting and rendering fallen stock, are performing the sampling of brains from fallen stock from most parts of Norway on behalf of the NFSA. The employees involved, mainly truck drivers, have been trained in the sampling by attending courses arranged by the NFSA.

The Norwegian Veterinary Institute (NVI) is performing the laboratory examinations and the reporting of the results.

## Aims

Scrapie surveillance is included in the Norwegian scrapie control programme in agreement with the requirements in the EU regulations.

## Materials and methods

In 2023, the surveillance programme was performed according to the European Union Regulations, Regulation (EC) No. 999/2001 Annex III, with amendments and included examination of the following categories of small ruminants:

- all small ruminants with clinical signs consistent with scrapie, irrespective of age.
- imported small ruminants, irrespective of age.
- dead, killed or slaughtered small ruminants older than 18 months from herds with restrictions and their contact flocks.
- 10,000 sheep older than 18 months, which had died or been killed on the farm (fallen stock).
- 10,000 randomly sampled healthy sheep older than 18 months slaughtered for human consumption.
- 500 goats older than 18 months which had died or been killed on the farm (fallen stock).

## Animals with clinical signs consistent with scrapie

When the sheep- and goat farmers recognized sheep or goats with clinical signs consistent with scrapie, they were responsible for notifying the case to the NFSA.

If indicated, the animals were subject to either post mortem examination at a laboratory, or formalin-fixed and unfixed brain halves and medial retropharyngeal lymph nodes were submitted for laboratory examination. All the samples were examined at the NVI.

## Surveillance of fallen stock

The sheep and goat farmers were responsible for reporting to the NFSA small ruminants older than 18 months that died or were killed on the farm due to disease. Employees from Biosirk collected the samples, which consisted of unfixed medulla oblongata and a small part of the cerebellum and midbrain obtained through the foramen magnum using a plastic spoon specially designed for the purpose. The samples were examined at the NVI.

## Abattoir surveillance

Brain samples from healthy sheep older than 18 months were collected by the NFSA. The sheep samples were collected at 22 abattoirs, which process all the commercially slaughtered sheep in Norway.

To ensure an appropriate distribution of the samples, the inspectors at the local NFSA were responsible for the sampling to be representative for each region and season. The sample selection should be designed to avoid overrepresentation of any group as regards to the origin, species, age, breed, production type or any other characteristic.

The brain samples consisted of medulla oblongata, and often a small part of the cerebellum and midbrain, obtained through the foramen magnum using the specially designed plastic spoon. The samples were examined at the NVI.

## Laboratory examination procedures

A rapid test (TeSeE<sup>®</sup> SAP ELISA, Bio-Rad or the HerdChek BSE-Scrapie Ag test, IDEXX) for detection of prions (PrP<sup>Sc</sup>) was performed for all submitted samples on a pooled brain tissue sample of obex and cerebellum when both structures were available or on the obex only when cerebellum was missing. In case of inconclusive or positive result, a western blot analysis (TeSeE<sup>®</sup> Western Blot, Bio-Rad) was used as confirmative test. The differentiation between classical scrapie and Nor98 scrapie was based on the Western blot profile.

## PrP genotyping

PrP genotyping was performed on all scrapie positive sheep. Genotyping of scrapie positive sheep was performed on unfixed brain samples at NVI. Genomic DNA was isolated using the DNeasy Blood & Tissue Kit (QIAGEN). Polymorphisms in the PrP gene were detected through Sanger sequencing of a PCR-generated product covering the entire coding region of the PRNP (forward primer 5' CAGTCATTCATTATGCTGCAGAC; reverse primer 5' CTATCCTACTATGAGAAAAATGAG).

From 2004 to 2016 genotyping of approximately 600 brain samples from healthy slaughtered sheep were performed each year to get an estimate of the PrP genotypes in the normal Norwegian sheep population.

## Prevalence

The scrapie prevalences in the fallen stock and abattoir populations were estimated assuming an exact binominal distribution.

# Results

## Sheep

Nor98 scrapie was diagnosed in seven sheep from seven flocks. Six cases were identified in fallen stock, while one case was an apparently healthy animal slaughtered for human consumption (Table 1). Since 1981, scrapie has been diagnosed in 278 different sheep flocks of which 221 have been flocks with Nor98 scrapie cases.

The individual age was registered, and the prion protein genotype examined for all scrapie cases (Table 2). All sheep, except one, had PrP genotypes with at least one allele with polymorphisms at codon 141 (AF<sub>141</sub>RQ) and/or codon 154 (AHQ). One sheep had the genotype ARQ/ARQ.

In total, 19,218 samples from sheep were received. Of these, 30 samples were unsuitable for examination. The numbers of animals examined within each category are presented in Table 1. The prevalence of Nor98 scrapie in the fallen stock of sheep was estimated to 0.06% [0.02-0.13%], (95% confidence interval, CI), and the prevalence of Nor98 scrapie in sheep slaughtered for human consumption was estimated to 0.01% [0.0003-0.06%], (95% CI) (Figure 2).

For 76 (0.4%) examined samples (40 healthy slaughtered and 35 fallen stock), the flock of origin was not reported. All the samples with unidentified origin of the flock were negative. In the event of a positive sample from slaughtered animals, the flock identity can be traced using the carcass number. The remaining 19,112 samples were collected from carcasses originating in 6,024 different sheep flocks. The mean number of animals tested per flock was 3.2 (range 1-82), sheep slaughtered for human consumption in restricted flocks due to scrapie are excluded. From 856 flocks, more than five animals were tested.

## Goat

Scrapie was not detected in goats in 2023. In total, 569 samples from goats were received. For 24 of these, the flock of origin was not reported (one healthy slaughtered, and 23 fallen stock). The numbers of animals examined within each category are presented in Table 1.

The collected samples originated from 210 different herds. The mean number of animals tested per herd was 2.6 (range 1-17). From 18 herds, more than five animals were tested.


*Table 1: Brain samples from sheep and goats submitted for examination for scrapie in 2023.*

Reason for submission to the laboratory	Number of samples			
	Total	Positive	Negative	Rejected
<b>Sheep - total</b>	<b>19,218</b>	<b>7</b>	<b>19,181</b>	<b>30</b>
Animals with clinical signs consistent with scrapie	8	0	8	0
Fallen stock	9,950	6	9,915	29
Healthy slaughtered animals	9,260	1	9,258	1
Animals examined due to scrapie restricted flocks	0	0	0	0
Imported animals	0	0	0	0
<b>Goats - total</b>	<b>569</b>	<b>0</b>	<b>568</b>	<b>1</b>
Animals with clinical signs consistent with scrapie	0	0	0	0
Fallen stock	556	0	555	1
Healthy slaughtered animals	13	0	13	0
Animals examined due to scrapie restricted flocks	0	0	0	0
Imported animals	0	0	0	0

*Table 2: Year of birth, reason for submission to laboratory examination, breed, prion protein genotype and type of scrapie of the scrapie cases detected in 2023.*

Case no	Year of birth	Reason for submission to laboratory examination <sup>1</sup>	Prion Protein Genotype	Scrapie type
1	2017	Fallen stock	AF <sub>141</sub> RQ/ARR	Nor98
2	2018	Healthy slaughtered animal	AF <sub>141</sub> RQ/AHQ	Nor98
3	2012	Fallen stock	AHQ/ARR	Nor98
4	2018	Fallen stock	AHQ/ARR	Nor98
5	2012	Fallen stock	AHQ/AHQ	Nor98
6	2015	Fallen stock	AHQ/ARQ	Nor98
7	2014	Fallen stock	ARQ/ARQ	Nor98

<sup>1</sup>The categories are: Healthy slaughtered animals, animals killed under scrapie eradication measures, suspect clinical signs consistent with scrapie including animals showing clinical signs at ante-mortem inspection, fallen stock (monitoring of fallen stock including animals examined because of other diseases than scrapie).



**Figure 2:** Box and whiskers plot of the prevalence of Nor98 scrapie from 2002 to 2023 in fallen stock (left) and slaughtered animals (right). The boxes represent the 25% to 75% quartiles and the whiskers the 2.5% and 97.5% exact binomial confidence intervals.

## Discussion

Nor98 scrapie was diagnosed in seven sheep, originating from seven different flocks. The age and genotypes of these sheep were in accordance with the previous experience of Nor98 scrapie (5). All cases, except one, had at least one of the alleles AF<sub>141</sub>RQ or AHQ that previously have been found to be associated with Nor98 scrapie (3).

The sheep positive for Nor98 scrapie were between five and eleven years old, which are in agreement with the result from previous years with the mean age being seven years (Table 2).

The Nor98 scrapie cases detected in 2023 were located in four different counties. In all of them, the disease had previously been diagnosed. Nor98 scrapie cases have been found in most parts of Norway and in all counties except Oslo, the capital of Norway. In contrast, the classical form of scrapie, has been detected only in the western part of Norway (two counties) and in Nordland County. From 2020, the Norwegian 19 counties were merged into 11 counties, and all numbers refer to the counties as of 2020.

The prevalence estimates of Nor98 scrapie in fallen stock and in sheep slaughtered for human consumption have varied during 2002-2023; however most estimates have been within the confidence intervals (Figure 2) (4). The results from the surveillance programmes indicate that the prevalence of Nor98 scrapie in the sheep population has not changed since the start of the programme.


Scrapie was not detected in goats in 2023. The first and only Nor98 scrapie case in goats in Norway was diagnosed in 2006 and originated from a county with a large goat population. Both classical and atypical scrapie in goats has been diagnosed in several countries in Europe (5).

## Acknowledgments

The authors thank all the persons who have contributed in sampling and the technical staff from the NVI for performing the analyses with excellence.

## References

1. Alvseike KR, Melkild I, Thorud K. Scrapie control at the national level: the Norwegian example. In: Hörnlimann B, Riesner D, Kretzschmar H (editors). Prions in humans and animals. Berlin: de Gruyter; 2006. p. 648-53.
2. Benestad SL, Sarradin P, Thu B, Schönheit J, Tranulis MA, Bratberg B. Cases of scrapie with unusual features in Norway and designation of a new type, Nor98. *Vet Rec.* 2003; 153: 202-8.
3. Moum T, Olsaker I, Hopp P, Moldal T, Valheim M, Moum T, Benestad SL. Polymorphisms at codons 141 and 154 in the ovine prion protein gene are associated with scrapie Nor98 cases. *J Gen Virol.* 2005; 86: 231-5
4. Mork, Jorunn, Benestad, Sylvie Lafond, Austbø, Lars, Moldal, Torfinn, Hopp, Petter. The surveillance programme for scrapie in Norway 2022. Surveillance program report. Veterinærinstituttet 2023. © Norwegian Veterinary Institute
5. Benestad SL, Arsac JN, Goldmann W, Nøremark M. Atypical/Nor98 scrapie: properties of the agent, genetics, and epidemiology. *Vet Res.* 2008; 39: 19.



*Scientifically ambitious, forward-looking  
and collaborative- for one health!*



**Veterinærinstituttet**  
Norwegian Veterinary Institute

Ås

Trondheim

Sandnes

Bergen

Harstad

Tromsø

postmottak@vetinst.no  
www.vetinst.no